



API - Canton

Attachment Parenting International

October 2006 Newsletter

A Well-Kept Secret

Breastfeeding's Benefits to Mothers

Alicia Dermer, MD, IBCLC
Old Bridge NJ USA

From: NEW BEGINNINGS, Vol. 18 No. 4, July-August 2001, p. 124-127

Very few people are unaware of the benefits of breastfeeding for babies, but the many benefits to the mother are often overlooked or even unknown. From the effect of oxytocin on the uterus to the warm emotional gains, breastfeeding gives a mother many reasons to be pleased with her choice. These documented effects are outlined in this excerpt from Breastfeeding Annual International 2001, a recently published anthology which was edited by Dia Michels, co-author of the classic breastfeeding advocacy book, *Milk, Money, and Madness*. Both books are available from LLLI.

One of the best-kept secrets about breastfeeding is that it's as healthy for mothers as for babies. Not only does lactation continue the natural physiologic process begun with conception and pregnancy, but it provides many short and long-term health benefits. These issues are rarely emphasized in prenatal counseling by health care professionals and all but ignored in popular parenting literature. Let's look at all the benefits breastfeeding provides mothers and speculate as to why so few are finding out about them.

Physiologic Effects of Breastfeeding

Immediately after birth, the repeated suckling of the baby releases oxytocin from the mother's pituitary gland. This hormone not only signals the breasts to release milk to the baby (this is known as the milk ejection reflex, or "let-down"), but simultaneously produces contractions in the uterus. The resulting contractions prevent postpartum hemorrhage and promote uterine involution (the return to a nonpregnant state).

Bottle-feeding mothers frequently receive synthetic oxytocin at birth through an intravenous line, but for the next few days, while they are at highest risk of postpartum hemorrhage, they are on their own. As long as a mother breastfeeds without substituting formula, foods, or pacifiers for feedings at the breast, the return of her menstrual periods is delayed (Lawrence and Lawrence 1999). Unlike bottle-feeding mothers, who typically get their periods back within six to eight weeks, breastfeeding mothers can often stay amenorrheic for several months. This condition has the important benefit of conserving iron in the mother's body and often provides natural spacing of pregnancies.

(con't on p.2)

API-Canton meets the second Tuesday morning of every month at 9:30. We are currently meeting at The Breastfeeding Center in Massillon.

Ideal #3: Breastfeed your baby – Breastfeeding meets baby's need for optimum nutrition and physical contact. Breastfeeding has many benefits to the baby, mother, and society and is the most natural way to meet so many of your baby's physical needs. While breastfeeding is the ideal way to feed a baby, parents who are not breastfeeding can still practice Attachment Parenting. We encourage parents who bottle-feed to use "breastfeeding" behaviors. In other words, hold your baby when feeding, talk to your baby and change positions during the feeding. Avoid the temptation to prop the baby's bottle since your baby will greatly benefit from your touch and holding.

* Advantages to mother and family:

1. Saves money – enough in one year to buy a major appliance
2. Saves time – no formula to prepare or bottles to wash
3. Convenient for home or travel
4. Triggers mothering hormones that promote attachment behaviors and calms mother
5. Helps mother can get more rest
6. Helps protect mother against breast cancer

* Advantages to baby:

1. Biologically designed for the human infant, contains needed nutrients in the proper amounts, digests easily
2. Gives immunity to certain diseases and viruses
3. Protects against some cancers, according to the newest research
4. Keeps baby close to mother and provides comfort
5. Helps strengthen jaws, eyes and formation of teeth
6. Less likely to develop allergies

* Avoid clock/calendar parenting. Follow your baby's cues rather than the clock or calendar.

* Weaning is a mutual process determined by the readiness of the baby and mother ("Cooperative weaning"). The World Health Organization (W.H.O.) code recommends breastfeeding until at least two years of age or beyond.

* If bottle-feeding, use breastfeeding behaviors:

1. Hold your infant when feeding, never prop the bottle
2. Make good eye contact at those times when your baby is alert and interested
3. Switch positions from one side to another; this helps strengthen the baby's eyes
4. Talk lovingly to your baby at feeding times

Membership Corner

NOW HIRING!

We are looking for co-leaders to help lead meetings. Requirements include: membership to API, having parented AP with one child for at least one year, and attendance to prior API meetings. We are also in need of a group treasurer, activities coordinator, enrichment meetings leader, fundraising coordinator, and greeter. In order to hold a group position, you must be a member. Please see Traci or Amy if you are interested.

Membership is not required to attend our meetings but your yearly dues are important to promote and sustain the activities of our group and support the work of API's national headquarters. With a yearly \$35 membership fee you are not only insuring continued education and support for yourselves and other families within your community but you will receive four quarterly issues of API news, have access to our book and sling lending libraries, have access to a research database and become a part of an international network of parents, professionals, and organizations who support attachment parenting. If you are interested in making flexible payments, please see us after the meeting, or by phone or e-mail.

Thank you to the following families who are supporting API through membership:

The Singree Family
 The Dunnerstick Family
 The Keoshian Family
 The Corey/Hall Family
 The Kemp Family

Welcome New Members:

The Mullane Family

(con't from p.1)

The amount of iron a mother's body uses in milk production is much less than the amount she would lose from menstrual bleeding. The net effect is a decreased risk of iron-deficiency anemia in the breastfeeding mother as compared with her formula-feeding counterpart. The longer the mother nurses and keeps her periods at bay, the stronger this effect (Institute of Medicine 1991).

As for fertility, the lactational amenorrhea method (LAM) is a well-documented contraceptive method, with 98 to 99 percent prevention of pregnancy in the first six months. The natural child-spacing achieved through LAM ensures the optimal survival of each child, and the physical recovery of the mother between pregnancies. In contrast, the bottle-feeding mother needs to start contraception within six weeks of the birth (Kennedy 1989).

Long-Term Benefits of Breastfeeding

It is now becoming clear that breastfeeding provides mothers with more than just short-term benefits in the early period after birth.

A number of studies have shown other potential health advantages that mothers can enjoy through breastfeeding. These include optimal metabolic profiles, reduced risk of various cancers, and psychological benefits.

Production of milk is an active metabolic process, requiring the use of 200 to 500 calories per day, on average. To use up this many calories, a bottlefeeding mother would have to swim at least 30 laps in a pool or bicycle uphill for an hour daily. Clearly, breastfeeding mothers have an edge on losing weight gained during pregnancy. Studies have confirmed that nonbreastfeeding mothers lose less weight and don't keep it off as well as breastfeeding mothers (Brewer 1989).

The above finding is particularly important for mothers who have had diabetes during their pregnancies. After birth, mothers with a history of gestational diabetes who breastfeed have lower blood sugars than nonbreastfeeding mothers (Kjos 1993). For these women who are already at increased risk of developing diabetes, the optimal weight loss from breastfeeding may translate into a decreased risk of diabetes in later life.

Women with Type I diabetes prior to their pregnancies tend to need less insulin while they breastfeed due to their reduced sugar levels. Breastfeeding mothers tend to have a high HDL cholesterol (Oyer 1989). The optimal weight loss, improved blood sugar control, and good cholesterol profile provided by breastfeeding may ultimately pay off with a lower risk of heart problems. This is especially important since heart attacks are the leading cause of death in women.

Another important element used in producing milk is calcium. Because women lose calcium while lactating, some health professionals have mistakenly assumed an increased risk of osteoporosis for women who breastfeed. However, current studies show that after weaning their children, breastfeeding mothers' bone density returns to pre-pregnancy or even higher levels (Sowers 1995). In the longterm, lactation may actually result in stronger bones and reduced risk of osteoporosis. In fact, recent studies have confirmed that women who did not breastfeed have a higher risk of hip fractures after menopause (Cummings 1993).

(con't on p.5)

Contacts:

API-Canton Co-Leaders:

Traci Singree - sunfairie@msn.com


Amy Dunnerstick - apmammaof2@yahoo.com

Secretary & Newsletter Info:

Dedra Keoshian - dedrakeo@yahoo.com

Book & Sling Librarian:

Jennifer Hall - halljene@muc.edu



Websites:

API-Canton Home Page:

<http://www.orgsites.com/oh/apicantoncircle/index.html>

API-Canton Discussion Forum:

<http://apicanton.org/index.php>

The Breastfeeding Center:

<http://www.thebreastfeedingcenter.com>
2800 Lincoln Way East (front door faces Elizabeth Ave)
Massillon, OH
330-837-0220

The Breastfeeding Center has been providing counseling and education to breastfeeding families since 1987. Our goal is to help you achieve a satisfying breastfeeding experience. We sell and rent quality breastpumps, nursing bras and other breastfeeding products. Instructions and demonstrations are provided with every purchase at no charge by Elizabeth Studer, RN, MSN, IBCLC, our board certified lactation consultant. Personalized nursing bra fittings are available with a wide selection of styles, colors, and sizes. We specialize in the hard to fit mother with a size range from 32A to 50K. Please call to set up an appointment.



PLEASE
remember to
return any
slings or

books that you have borrowed. Remember, slings are for RENT at \$3.00 per month. Both slings and books should be returned as soon as possible so that other members can borrow them.

Our Mother's Helper

We have a Mother's Helper that joins us to care for the older children while we chat, through games, stories, and crafts. If your child plays with her, we ask that you be prepared to donate at least \$1 per child for her valuable and playful time!

Upcoming Events

Oct. 8 – Safety Day at Walsh Farms
There will be med-flight, helicopters, fire trucks, sheriff, police cars, farm equipment and much much more.:
<http://walshfarms.net/>

Oct. 14 – The Enchanted Forest Walk

Non-breastfeeding mothers have been shown in numerous studies to have a higher risk of reproductive cancers. Ovarian and uterine cancers have been found to be more common in women who did not breastfeed. This may be due to the repeated ovulatory cycles and exposure to higher levels of estrogen from not breastfeeding. Although numerous studies have looked at the relationship between breastfeeding and breast cancer, the results have been conflicting. This is largely due to flaws in study design and lack of uniform definition of breastfeeding, resulting in difficulty comparing the data. (In some studies, breastfeeding has been defined as having breastfed at least once a day, while in others it is defined as exclusive breastfeeding, using no supplements or artificial nipples.) Despite this, it is now estimated that breastfeeding from six to 24 months throughout a mother's reproductive lifetime may reduce the risk of breast cancer by 11 to 25 percent (Lyde 1989; Newcomb 1994). This phenomenon may also be due to suppressed ovulation and low estrogen, but a local effect relating to the normal physiologic function of the breast may also be involved. This was suggested by a study in which mothers who traditionally breastfed on only one side had significantly higher rates of cancer in the unsuckled breast (Ing, Ho, and Petrakis 1977).

In two studies, there appeared to be an increase in flare-ups of rheumatoid arthritis in breastfeeding mothers (Jorgensen 1996; Brenna 1994). However, in another study, overall severity and mortality of rheumatoid arthritis was worse in women who had never breastfed (Brun, Nilson, and Kvale 1995). There have been no other studies showing any detrimental health effects to women from breastfeeding. Bottom line: Breastfeeding reduces risk factors for three of the most serious diseases for women-female cancers, heart disease, and osteoporosis-without any significant health risks.

Psychological Issues for Breastfeeding Mothers

How do you measure the peace of mind of having a healthy baby who is developing optimally? Where do you factor in the financial burden of formula prices and increased medical costs?

Public health agencies advocate for breastfeeding because of its well-documented health advantages to babies, but they fail to convey to individual mothers and families the potential emotional impact of this very crucial infant-feeding decision. In Western society, the decision about breast or bottle is still seen very much as a personal choice based on convenience. The potential stress of living with a child with recurrent illnesses, or the loss of the unique bond that comes from breastfeeding, are often omitted from the decision-making process.

There is much more to breastfeeding than the provision of optimal nutrition and protection from disease through mother's milk. Breastfeeding provides a unique interaction between mother and child, an automatic, skin-to-skin closeness and nurturing that bottle-feeding mothers have to work to replicate. The child's suckling at the breast produces a special hormonal milieu for the mother. Prolactin, the milk-making hormone, appears to produce a special calmness in mothers. Breastfeeding mothers have been shown to have a less intense response to adrenaline (Altemus 1995).

This calming effect is hard to measure in a society largely unsupportive of breastfeeding such as the United States, where breastfeeding beyond the early weeks is not the norm. Mothers who try to breastfeed in this climate often experience physical and emotional problems. These problems result from a lack of breastfeeding role models among family and friends, and are compounded by the easy availability of formula and a lack of access to knowledgeable and supportive health care professionals.

(con't on p.6)

Even if a mother overcomes physical problems, she may still encounter negative comments, such as "Are you still nursing?" or "Your milk may not be strong enough-why don't you add formula?" Or her employer may make it impossible for her to continue breastfeeding on returning to work. Or she may be harassed for breastfeeding in public. No wonder that few mothers get to fully experience the relaxing effects of breastfeeding.

New motherhood is a time fraught with emotion. The baby blues are common, often exacerbated by lack of support and a sense of isolation. The role of breastfeeding in postpartum emotional upheavals has not been well studied, but breastfeeding mothers with depression need treatment just as much as any other mother. Such women present a unique challenge to health care professionals. Since medications may pass into breast milk, many physicians believe the safest solution is to wean the child. However, in most cases of depression, women do better if they continue to breastfeed. Unfortunately, too often physicians insist that mothers wean their child in order to take antidepressant medicines.

A review of the literature, however, has demonstrated that several antidepressants pose minimal, if any, risk to the nursing child. A mother who feels that her nursing relationship with her child is the only thing going right in her life can now continue to breastfeed while receiving appropriate medications for her depression.

Why Don't More People Know How Good Breastfeeding Is?

Clearly, breastfeeding is good for mothers both physically and emotionally. And yet, many mothers decide to breastfeed based solely on the benefits to the baby. Breastfeeding in the context of a bottle-feeding society tends to be perceived as inconvenient and uncomfortable.

Often, mothers see breastfeeding as martyrdom to be endured for their baby's health. If they stop early, they may feel guilty about depriving the baby of some health benefits, but their guilt is often soothed by well-meaning people who reassure them that "The baby will do just as well on formula." Perhaps if they knew that continuing to breastfeed is also good for their own health, some mothers might be less likely to quit when they run into problems.

Many mothers are not being told how good breastfeeding is for their health. Whether out of ignorance or due to the influence of the artificial baby milk industry, many health care providers fail to inform mothers of the facts. It's time for this well-kept secret to come out. As word spreads about these little-known facts, more mothers will not merely choose to breastfeed briefly to provide early disease protection for their baby, but will continue to breastfeed, providing optimal outcomes both for their children and for themselves.

Alicia Dermer, MD, IBCLC, is Clinical Associate Professor in the Department of Family Medicine at the University of Medicine and Dentistry of New Jersey-Robert Wood Johnson Medical School in New Brunswick, New Jersey. She has a special interest in wellness and health promotion. As part of this interest, she has gained expertise in breastfeeding education and promotion. She successfully sat for the certifying examination of the International Board of Lactation Examiners in 1995. She lectures extensively on the subject of lactation, is actively involved in health care professional and lay education about breastfeeding, and has several publications on the subject.